

# The SPoRT Center – Infusing NASA Technology Into NWS WFOs

# Dr. Gary Jedlovec, NASA / MSFC Earth Science Office

Mission of the SPoRT Center: Apply NASA measurement systems and unique Earth science research to improve the accuracy of short-term (0-24 hr) weather prediction at the regional and local scale

SPoRT website

http://weather.msfc.nasa.gov/sport
Collaborative Partner Blog
http://www.nsstc.uah.edu/sportblog/



SPoRT - Short-term Prediction and Research Transition





# SPoRT - A NASA Test Bed and More!

#### SPoRT is an end-to-end research to operations activity!

- explore cutting edge research relevant to operational weather forecasting
- work with end users to match forecast problems to capabilities / data

# Demonstrate feasibility of new data / capability to address forecast problem in a simulated operational environment – this is a test bed function

- real-time data and products timely dissemination capability
- display in operational system visualization, interoperability with other products
- assessment and impact on forecast problem
- end user involvement critical to success

### Transition of experimental data to operational environment

- mechanics of transition AWIPS / AWIPS2 and other DSS
- training and product impact and assessment end user involvement








# SPoRT Products to End Users

# Active collaborations with 14 WFOs – mainly in Southern Region

- Suite of over 30 products integrated into AWIPS
- MODIS, AMSR-E, special GOES products, and total lightning data
- WRF forecasts (from HWT)
- LIS and local surface analyses
- High resolution SST analyses for WRF EMS users

### **Expanding to Other Regions**

- Requested by SSD chiefs March 2010
- Project-based expansion with existing product suite
- New regional forecast problems







## Forecast Problems / Test Bed Collaborations

#### **Southern Region Forecast Problems**

- timing and location of severe weather
- morning minimum temperatures (local variations)
- detection and monitoring of fog, smoke, fires
- coastal weather processes (sea breeze convection / temperatures)
- development / movement of off-shore precipitation processes – tropical systems
- gap filler in data void regions atmospheric rivers of moisture



#### **NOAA Test Bed Collaboration**

- HWT forecast program
  - NASA SST data and configurations of NSSL WRF forecasts forecasts to WFOs
  - Lightning forecasts (derived from microphysics in WRF forecasts)
- GOES-R Proving Ground
  - total lightning data for multiple ground based networks
  - pseudo GLM product for forecaster evaluation





# **MODIS Applications**

#### **Relevance to NWS Forecasters**


- ■Infuses NASA EOS data into the short-term forecast process.
- Serves as an introduction to future polar and geostationary products

# Unique SPoRT/NASA Contribution

■High resolution visible imagery, false color products, detection of fog in small terrain features, and hot spot detection.

#### **Future Work**

- Additional products to address specific forecast problems
- ■Blend MODIS and GOES imagery as MODIS becomes available, allowing for a "hybrid" product







# Operational Uses of Lightning Mapping Array

#### **Relevance to NWS Forecasters**


■Some NWS WFOs have access to lightning mapping array data, others will have future access through the GLM on GOES-R.

#### **Unique NASA/SPoRT Contributions**

- ■Working to obtain total lightning data from a variety of CONUS sources (Alabama, DC, Florida, New Mexico, Oklahoma)
- ■Incorporation of total lightning data within AWIPS/AWIPS 2 for overlay with traditional radar and other analyses
- ■Translate total lightning trends into severe weather precursors

#### **Future Work**

■ Exploring new tools available within AWIPS or AWIPS 2 that permit tracking of flash rates by cells for use with lightning trend algorithms





transitioning unique NASA data and research technologies to operation



# 3D Analysis Product Using Hyperspectral Sounder Profiles

#### **Relevance to NWS Forecasters**

- Profiles from hyperspectral sounders (e.g. AIRS and IASI) provide rawinsonde-quality observations to fill spatial and temporal data voids
- Temperature and moisture profiles can be used to improve diagnostic analysis, atmospheric rivers, and moisture return forecasts

#### **Unique SPoRT/NASA Contribution**


- Configured WRF-Var analysis system to assimilate AIRS thermodynamic profiles
- Rigorous testing of analysis system on 37 case study days from winter 2007

#### **Future Work**

- Incorporate IASI profiles into analysis
- Generate and distribute real-time product to WFOs

Improved 6-hr cumulative precipitation from improved CAPE and 1000 hPa dew point









# Sea and Lake Surface Temperature Composites

#### **Relevance to NWS Forecasters**


 Observations of sea and lake surface temperatures provide high resolution detail for coastal and marine forecasts

#### **Unique SPoRT/NASA Contributions**

- high resolution (1km) MODIS and AMSR-E composites – 4x daily
- Provides coverage throughout North
   America and the tropics, depicts coastal and
   Gulf Stream gradients
- ■Developed a unique application over the Great Lakes that includes lake ice analyses from the NOAA Great Lakes Environmental Research Laboratory (GLERL)

#### **Future Work**

■Examine sensitivities of WRF-EMS forecasts to initialization with SPoRT SSTs and Great Lakes temperature composites.







# Applications of the Land Information System (LIS)

#### **Relevance to NWS Forecasters**


- ■NASA land surface modeling and assimilation system provides high-resolution soil data for NWP
- ■Reduces the diurnal 2-m temperature bias in model forecasts and improves QPF for convective events

#### **Unique SPoRT/NASA Contributions**

- ■SPoRT maintains a near real-time run of the NASA LIS, providing output over the southeastern U.S.
- ■High resolution (3 km) output available for WRF-EMS initialization, currently used by NWS Miami, FL

#### **Future Work**

- Incorporate MODIS vegetation data to improve representation of the land surface
- ■Provide output LIS fields in AWIPS/2 for diagnostic evaluation at partner WFOs





# ontributions to WRF Environmental Modeling System

#### **Relevance to NWS Forecasters**


■The WRF-EMS is used within several NWS WFOs to provide local, high resolution forecasts for prediction and training.

#### **Unique SPoRT/NASA Contributions**


- ■The SPoRT SST product is the *current default* within the WRF-EMS, effective V3.1, and real-time composites are distributed four times daily in GRIB format.
- ■SPoRT provides LIS initialization fields for the WRF-EMS, currently available over the southeastern United States.

#### **Future Work**

- ■Examine impact of Great Lakes temperatures on lake effect weather.
- ■Broaden the LIS domain to include a broader sample of the United States.











# **Use of Experimental Data in Operations**

## **SPoRT Best Practices**

- End user interactions
  - understand needs, the operational environment, and forecast constraints – "live in their world" for awhile
  - involve end user in entire process
  - site visits, coordination calls (GoToMeeting), collaborative blogs, collaborative workshops
- Integrate into user DSS (AWIPS, AWIPS2, etc.)
- Produce user training on products
- Document successes
- Communicate results in various forums








# **External Communication and Outreach**

Outreach via web, blog, Facebook, publications - more visibility to external community

- Peer-reviewed publications, quarterly and biennial reports
- Revamped SPoRT website, Wide World of SPoRT blog in (over 100 posts since March 2009), >150 Facebook fans (mainly NWS)













# **Future Direction**

Continue to demonstrate utility of current and future NASA observations

# **Decision support systems**

- •AWIPS / AWIPS2
- •Google Earth

# JPSS (NPP / NPOESS)

•VIIRS and CrIMSS data and products to WFOs

# **Expand partners and end users**

- Work more collaboratively with regional centers and other testbeds
- Expand to WFOs in other regions

